

REVISED VERSION

(Note: A previous version made grandiose unsustainable claims.)

An axiomatic account of space sufficient to provide formal proofs of spatial relationships

ABSTRACT

A definition is given of *seriate sets* as being sets constituted out of structured collections of objects which are recursively internally self-similar. Fundamental (geometrical) objects of Dimension N are conceived to be constituted out of seriate sets of fundamental objects of Dimension $N-1$, starting with points assigned to Dimension 0. Syntactical rules to enable such objects to be systematically named and combined, are set out. A series of formal proofs of theorems relative to objects of Dimensions 1 and 2 are worked through. A proof that four colours are sufficient to colour any five area map is given in illustration.

INTRODUCTION

This paper describes an axiomatic calculus which can be interpreted as descriptive of multidimensional *a-mensurate* space: that is space which can be conceived of as existing even before the notion of mensuration has been arrived at. This calculus is founded on the idea of a *seriate set*. Seriate sets are understood to be sets possessing recursive structural internal self-similarity. Seriate sets of points are interpreted as lines, seriate sets of lines as areas, and so on, to establish a multidimensional system.

The first section of the paper, entitled *Foundations*, sets out a purely abstract definition of seriate sets, that is without any reference to geometrical objects. The section following, headed *Dimension 1*, is concerned with seriate sets of points as constituting *fundamental lines*. The first two theorems here deal with the *external* and *internal* self-similarity of fundamental lines: that is with the capacity of two such lines to conjoin to form an exactly equivalent fundamental line, and with the ability of any single fundamental line to be divided by any (internal) point into two exactly equivalent fundamental lines. The remainder of this section includes theorems to the effect that any non end-point of a fundamental line is between its end-points, while neither of those end-points is ever between any other two points of that line; and that one, and one only, of three (different) points within a fundamental line is between the other two, for example.

The next section is concerned with objects of Dimension 2. Fundamental characteristics of seriate sets constituted out of fundamental lines are demonstrated in a number of theorems. Such seriate sets are then interpreted as constituting fundamental areas. Fundamental areas are shown to be externally and internally self-similar analogously to fundamental lines. These results then enable a proof to be given that four colours are sufficient for any five country map where the countries are all fundamental areas within a given fundamental area.

The paper is divided into separate sections each consisting of a formal Definition, or Convention, or Theorem, or the like, typically followed by a short section of commentary. A few brief general

notes are added at the end. An appendix is added giving more rigorous proofs of the first two theorems of the main text, which there, for the sake of brevity and the reader's convenience, are given in a more intuitive presentation. Illustrative diagrams, purely as an aid to comprehension, are added.

FOUNDATIONS

DEFINITION (D) 0.1: THE AXIOM OF EXISTENCE

The existence of any object can be asserted if that existence does not entail "not A" where 'A' stands for any proposition entailed by any characteristic of any array of which the given object is asserted to be a part; and can be negated if it does entail "not A".

An ARRAY is any collection of well-formed objects of the calculus. The assertion of the existence of an object can be imagined to be roughly analogous to drawing a line on a sheet of paper: we can draw any specified line we like so long as this does not offend against something already drawn. For example we can draw a line representing the diagonal of a square which does not intersect any other line - so long as the other diagonal has not previously been drawn.

DEFINITION 0.2: SERIATE SETS

A set of more than two like objects constitutes a seriate set $S!$, if and only if:

- (a) no part of any object which is a member of $S!$ is a part of any other object which is also a member of $S!$;
- (b) the membership of $S!$, includes two, and only two, objects, termed "E" objects of $S!$, while all the other members of $S!$ are termed "I" objects of $S!$, in such wise that given the existence of $S!$ and given any "I" object within $S!$, the existence can be asserted of a set of seriate sets $*S$ which fulfils the following conditions:
 - (i) the set of all the members of all the seriate sets of $*S$ is identical one-to-one with all the set of all the members of the given seriate set $S!$;
 - (ii) no member of any given seriate set of $*S$ is a member of any other seriate set of $*S$ if it is not an "E" object of that seriate set;
 - (iii) two, and only two, of all the "E" objects of all the seriate sets of $*S$, are members of no other seriate set of $*S$, and are identical to the two "E" objects of $S!$;
 - (iv) every other "E" object of any seriate set of $*S$ is a member of two and only two seriate sets of $*S$, and the given "I" object is identical to such an "E" object;
 - (v) no subset of sets of $*S$ is such that each of the "E" objects of every member of that subset is a member of another seriate set of that subset.

See diagrams (at end) for the illustration of a seriate set of points in accord with the above. LIKE objects are all of the same dimension, as are points or lines. [Dimension is formally defined p. 8.] Variants on the form of $S!$ such as $S^6!$ and $S^N!$ are all names of seriate sets; and variants on the form of $*S$ such as $*S^2$ and $*S^J$ are all names of (simple) sets, consisting of unstructured aggregations of objects. An ELEMENT of an object is any part of that object which CONSTITUTES that object, that is, is a MEMBER of the (seriate) set which determines the characteristics of the object.

DEFINITION 0.3: THE AXIOM OF IDENTITY

One set is IDENTICAL to another if it is not possible to distinguish the membership of one set from the membership of the other.

To DISTINGUISH here refers to the unitary process by which objects are isolated/identified/named.

DEFINITION 0.4: THE AXIOM OF STABILITY

If, given a seriate set $S!$, the existence of a seriate set $S^N!$ within $S!$ can be asserted, then the existence of any other seriate set within $S!$ cannot be asserted, which set has same "E" objects as $S^N!$ and is constituted by a set of objects which is not identical to the set of objects which constitutes $S^N!$.

Anything GIVEN it is to be understood to have had its existence asserted. An object or set is WITHIN another object or set if every element or member of the first object or set is likewise an element or member of the second object or set. D. 0.2 above ensures that a unique serial order can be established between all the objects of a seriate set. It has nothing to say however about any other possible order those same objects might (?) have. This axiom ensures that once one serial order in a seriate set has been established no other conflicting order can be asserted to exist.

DIMENSION 1

INTERPRETATION (In.) 1.1

A seriate set of points constitutes a FUNDAMENTAL LINE; and the "E" objects of that seriate set are termed the END-POINTS of that line.

A POINT is formally defined in the familiar way as that which has no parts.

CONVENTIONS (Cn.) 1.1

- (i) The names of points take the form of unadorned upper case letters, e.g. 'A', 'B', 'N'.
- (ii) The names of fundamental lines take the form of AB where 'A' and 'B' are the end-points of the line. The objects - points - which constitute that line are referenced by means of a variable, always given in lower case, as for instance 'x', in the formulation AB(x).
- (iii) The use of round brackets containing a variable immediately following the name of an object, or set, is the general syntax for the assignment of a variable which references all and only, the elements of the named object or members of the given set. In the case of a fundamental line the name of a point may be included within those same brackets in addition to indicate that it is an element of the given line. E.g. AB(x,C) states 'C' is a point in the same line.
- (iv) Objects having different names are to be assumed non identical (the default condition) until the converse is stipulated or proved.
- (v) Symbols and syntax: '&', ' $\wedge$ ', stand for the logical operators 'and' and 'or'. ' $\neg$ ' is used for negation. ' $\implies$ ' and ' $\iff$ ' represent "if...then" and "if and only if"; ' $\longmapsto$ ' represents a one-to-one identity mapping, i.e. ' $x \longmapsto y$ ' means every object referenced by 'x' is identical one-to-one with an object referenced by 'y'; ' $\longleftrightarrow$ ' means this relationship is reciprocal indicating identity of reference. '+' indicates the addition of variables and '-' their subtraction where, and only where, the second variable named references a subset of the objects referenced by the

first. Where the mapping arrow, or the identity sign '=', is used in conjunction with the logical operators ' $\wedge$ ' or '&': ' $\wedge$ ' means that at least one of the arguments of that operator must have the relationship indicated, and '&' means both must have it. The meaning of any string of symbols in any case where ambiguity might arise is determined by assigning precedence to the leftmost member of any pair of operators.

(vi) In proofs, the writing of the bare unmodified name of an object represents the proposition that that object exists. Such propositions then take truth values. The unadorned name of an object can then appear in any argument place of a logical operators, always, it must be remembered, acting as surrogate for the proposition that the named object exists.

Sect. (iii): The reference of variables is confined to objects all of the same dimension. Sect. (v): $[P \iff Q]$ is more precisely defined as equivalent to $\{[P \implies Q] \& [Q \implies P]\}$. Operator precedence: $\{[A \& B] \mapsto [x \wedge y]\}$ means $\{[A \mapsto (x \wedge y)] \& [B \mapsto (x \wedge y)]\}$, and not $\{[(A \& B) \mapsto x] \wedge [(A \& B) \mapsto y]\}$. $\{a \mapsto (x \wedge y)\}$ means $\{[a \mapsto x] \wedge [a \mapsto y]\}$ and not that some of the objects referenced by 'a' map onto 'x' and/or some of them map onto 'y'. $\{P \mapsto \neg a\}$ is not allowed. The identity sign indicates identity between the objects referenced and not merely identity between signs. Thus $\{C = [A \wedge B]\}$ means that some point 'C' is identical to 'A' or 'B' - or possibly both - and not that 'A' or 'B' can be replaced by 'C' in any statement containing either.

THEOREM (Th.) 1.1

Given any two fundamental lines AB and BC which have no point in common except 'B', then the existence of a fundamental line AC constituted by all and only the points which constitute AB and BC can be asserted.

1. $\underline{AB}(x) \& \underline{BC}(y)$
2. $\neg \{[P \mapsto (x \& y)] \& \neg [P = B]\}$
3. $[T \mapsto x] \& \neg [T = (A \wedge B)]$
4. $\{\underline{AN}^1(n^1) \& \underline{N}^1\underline{N}^2(n^2) \& \underline{N}^2\underline{N}^3(n^3) \& \dots$
 $\dots \& \underline{N}^{K-1}\underline{N}^K(n^K) \& \underline{N}^K\underline{B}(n^{K+1})\}$
 $\& \{x \longleftrightarrow [n^1 + n^2 + n^3 \dots n^{K+1}]\}$
 $\& \{T = [N^1 \wedge N^2 \wedge N^3 \dots N^K]\}$ 1,3,D.0.2
5. $z \longleftrightarrow (x + y)$
6. $[\underline{AN}^1(n^1) \& \underline{N}^1\underline{N}^2(n^2) \& \dots$
 $\dots \& \underline{N}^{K-1}\underline{N}^K(n^K) \& \underline{N}^K\underline{B}(n^{K+1}) \& \underline{BC}(y)]\}$
 $\& \{z \longleftrightarrow [n^1 + n^2 + n^3 \dots n^{K+1} + y]\}$
 $\& \{T = [N^1 \wedge N^2 \wedge N^3 \dots N^K \wedge B]\}$ 1,4,5
7. $\underline{AC}(z)$ 2,6,lt.(T),Sym.D.0.2

L(LINE) 2: This simply states AB and BC have no point in common except 'B'. L. 3. That a point such as 'T', distinct from 'A' and 'B' can be asserted to map onto 'x' arises from the first clause of D.0.2, but such is

generally taken to be too obvious to need listing in the third column; further it is not generally though necessary to first assert the existence of such an object before asserting such a relationship, except where necessary for emphasis. This pattern holds throughout. L. 7: When the default condition of Cn.1.1(iv) is read as strongly as possible at L. 4 then given L. 2, D.0.2(b) clauses (ii)/(iv) hold for 'T' at L. 6. By iteration and similarity what follows for 'T' follows for all non end-points of AB and BC. The same holds for 'B' directly from L. 1/2. D.0.2(b)v in effect precludes any subset of fundamental lines of such a set as that of L. 4, forming a "separate loop". As it holds for each of AB and BC in isolation, if it were controverted when they are conjoined then they would have to have more than one point in common contravening L. 2. All the points of 'z', not being 'A' or 'C' then fulfil the requirements of a seriate set "I" object of AC(z). See Appendix for a more rigorous proof of this and the next theorem.

THEOREM 1.2

Given any non end-point within a fundamental line AB, then the existence of two fundamental lines AC and CB, within AB, constituted out of all and only the elements of AB, and having no point in common except C, can be asserted.

1. AB(C,x)
2. $\{\underline{AX}^1(x^1) \ \& \ \underline{X^1X^2}(x^2) \ \& \ \underline{X^2X^3}(x^3) \ \& \ \dots$
 $\dots \ \& \ \underline{X^{K-1}X^K}(x^K) \ \& \ \underline{X^KB}(x^{K+1})\}$
 $\& \ \{x \longleftrightarrow [x^1 + x^2 + x^3 \dots x^{K+1}]\}$
 $\& \ \{C = [X^1 \wedge X^2 \wedge X^3 \dots X^K]\}$ 1,D.0.2
3. $C = X^1$
4. AC(y¹) & [y¹ $\longleftrightarrow$ x¹] 2:3
5. CX²(x²) & X²X³(x³) & ... X^KB (x^{K+1}) 2:3
6. $z^1 \longleftrightarrow (x^2 + x^3 + \dots x^{K+1})$
7. CB(z¹) & [y¹ + z¹] $\longleftrightarrow$ x 5,6,2,Th.1.1,lt
8. $C = X^2$
9. AC(y²) & [y² $\longleftrightarrow$ (x¹ + x²)] 8,Sm. 5/7
10. CB(z²) & [z² $\longleftrightarrow$ (x³ + x⁴ + ... x^{K+1})] 8,Sm 5/7]
11. [y² + z²] $\longleftrightarrow$ x 2:9,10
12. AC(y) & CB(z) & [(y + z) $\longleftrightarrow$ x] &
 $\neg \{[P \longrightarrow (y \ \& \ z)] \ \& \ \neg [P = C]\}$ 2,lt.C(X^N)

L. 4: The colon separating the line number in column three indicates the embedding of an entailment. L.12: Iterating for 'C' identical to every 'X' in L. 2. D. 0.2(b) clause (v) obviously holds throughout. See Appendix for a more rigorous proof of this theorem.

THEOREM 1.3

Given any two points 'P' and 'Q' within a given fundamental line $\underline{AB}$, then the existence of a fundamental line $\underline{PQ}$ within $\underline{AB}$ together with two other fundamental lines within $\underline{AB}$ - either $\underline{AP}$ and $\underline{QB}$, or $\underline{AQ}$ and $\underline{PB}$ - can be asserted, such that no point within $\underline{AB}$, apart from 'P' and 'Q', is an element of both $\underline{PQ}$ and either of the those other two fundamental lines, which three lines exhaust $\underline{AB}$.

This follows directly from Th. 1.2 iterated.

INTERPRETATION 1.2: BETWEEN

A point 'B' is BETWEEN two other points 'A' and 'C', if and only if, within some given set of points, the existence of two fundamental lines $\underline{AB}$ and $\underline{BC}$ can be asserted, which have in common no other point but 'B'; and this between relationship is said to exist *within* the given set of points. It is symbolised $A/B/C(x)$ where 'x' references all the points in the given (underlying) set.

It is crucial that throughout one keeps firmly in mind that the "between relationship" is a relationship that exists only in respect of some underlying set.

THEOREM 1.4

Every non end-point of a given fundamental line set is between the end- points of the given line, within that line.

This follows directly from Th. 1.2 and In. 1.2.

THEOREM 1.5

One point of any three (different) points within a fundamental line is between the other two, within that line.

Apply Th. 1.3 to a pair of the given points and then Th. 1.2 to the third point, and In. 1.2 will hold in the case of one of those given points.

THEOREM 1.6

If one point is between two others within a given fundamental line, then neither of the others is between the other two, within that line.

1. $\underline{AB}(H,J,K,x)$
2. $H/J/K(x)$ 1,Th.1.5,Rn.
3. $\underline{HJ}(r) \ \& \ \underline{JK}(s) \ \& \ [(r \ \& \ s) \longmapsto x]$
 $\ \ \ \ \ \& \ \neg \{[P \longmapsto (r \ \& \ s)] \ \& \ \neg [P = J]\}$ 2,In.1.2
4. $\underline{HK}(t) \ \& \ [t \longleftrightarrow (r + s)]$ 3,Th.1.1
5. $J \longmapsto t$ 3,4
6. $J/H/K(x)$
7. $\underline{JH}(r^1) \ \& \ \underline{HK}(t^1) \ \& \ [(r^1 \ \& \ t^1) \longmapsto x]$

	$\& \neg \{[P \mapsto (r^1 \& t^1)] \& \neg [P = H]\}$	6,ln.1.2
8.	$\neg [J \mapsto t^1]$	7
9.	$\neg [t \longleftrightarrow t^1]$	5,8
10.	$\neg [H/J/K(x) \& J/H/K(x)]$	9,4,7,1,D.0.04

L. 2: Rename the points H,J,K so J is between the other two if it is not already so. It follows from this theorem that "between relationships" within a line commute to form a single ordering. This ordering can then be symbolised through an extrapolation of the basic "between syntax", as for example in A/B/C/D(x).

THEOREM 1.7

Given any pair of points within a given fundamental line then neither end-point of that line is between that pair of points, within that line.

Apply Th. 1.2 and this result then follows directly from Th. 1.4 and Th. 1.6, in both possible cases.

INTERPRETATION 1.3

If, and only if, a set of points constitutes a RING (set), then all and only, those same points, constitute two fundamental lines which have solely in common the end-points of each.

THEOREM 1.8

Given any pair of points each within a pair of fundamental lines which constitute a ring, then the existence of a pair of fundamental lines with those points as end-points can be asserted, which pair of lines constitutes a ring constituted out of all and only the points which constitute the original pair of lines.

1.	$\underline{AB}(a) \& \underline{BA}(b)$	
2.	$\neg \{[P \mapsto (a \& b)] \& \neg [P = (A \wedge B)]\}$	
3.	$[a + b] \longleftrightarrow z$	
4.	$[F \& G] \mapsto a]$	
5.	$\underline{AF}(c) \& \underline{FB}(d) \& [(c + d) \longleftrightarrow a]$	4Th.1.2
6.	$G \mapsto c$	4,5,Rn.
7.	$\underline{AG}(e) \& \underline{GF}(f) \& [(e + f) \longleftrightarrow c]$	6,Th.1.2
8.	$\underline{GF}(g) \& [g \longleftrightarrow (e + b + d)]$	7,1,2,5,Th.1.1
9.	$[f + g] \longleftrightarrow z$	8,7,5,3
10.	$\underline{GF}(f) \& \underline{GF}(g) \& [(f + g) \mapsto z]$	7/9
11.	QED 10,Sym.,It.	

L. 11: L. 4 assumes the two given points are within the same given line. By symmetry the same result holds for both given lines. If the two given points are each within a different one of the two given fundamental lines, then by judiciously selecting a pair of intermediate points a simple iteration of L.10 achieves the same result.

Corollary: Given any pair of points within a ring then the existence of a pair of lines with those points as end-points and no other points in common can be asserted constituted out of the same set of points as constitutes the ring.

Given the ring in the first place then the situation as described in L.1 must hold by In. 1.3 and hence, given this together with the main proof the same applies to every pair of points within the ring.

CONVENTION 1.2

The general form of the name for a ring is $X//.../X(x)$, where 'X' is any point in the ring and 'x' is a variable which references all (and only) the points within the set of points that constitutes the ring: the names of any points within the ring can be filled in between the slashes, or omitted, as required, in accord with the convention for the between relationship.

THEOREM 1.9

Every one of three points of a ring is between the other two within the set of the ring.

1. $A//.../A(x)$
2. $[B \ \& \ C] \mapsto x$
3. $\underline{AB}(a) \ \& \ \underline{BA}(b) \ \& \ [(a + b) \longleftrightarrow x]$
 $\ \ \ \ \ \& \ \neg \{[P \mapsto (a \ \& \ b)] \ \& \ \neg [P = (A \ \wedge \ B)]\}$ 1,2,Th.1.8,Cr.
4. $C \mapsto a$ Rn.
5. $\underline{AC}(c) \ \& \ \underline{CB}(d) \ \& \ [(c + d) \longleftrightarrow a]$
 $\ \ \ \ \ \& \ \neg \{[P \mapsto (c \ \& \ d)] \ \& \ \neg [P = C]\}$ 3,4,Th.1.2
6. $\underline{AC}(c) \ \& \ \underline{CB}(d) \ \& \ \underline{BA}(b) \ \& \ [(c \ \& \ d \ \& \ b) \mapsto x]$ 3,5
7. $\neg [P \mapsto (A \ \wedge \ B \ \wedge \ C)] \implies$
 $\ \ \ \ \ \neg \{P \mapsto [(c \ \& \ d) \ \wedge \ (c \ \& \ b) \ \wedge \ (d \ \& \ b)]\}$ 3,5
8. $A/C/B(x) \ \& \ C/B/A(x) \ \& \ B/A/C(x)$ 6,7,In.1.2.

L. 4: Reverse the names of the variables 'a' and 'b' if 'C' is not as indicated.

THEOREM 1.10

No set of points which forms a ring is within a fundamental line.

This follows directly from Th. 1.9 and Th. 1.6

DIMENSION 2

INTERPRETATION 2.1

A seriate set constituted out of objects of DIMENSION (N), and the object which that set constitutes, both belong to Dimension (N + 1). A point belongs to Dimension 0. The dimension of a seriate set is indicated by a subscript: a set of points, for example, is written $S_1!$ (the default condition so the subscript is often dropped in this case), and a seriate set of lines $S_2!$. When the subscript for dimension is used the superscripts differentiating different seriate sets are displaced to the end of the name so that $S_2!^N$, and $S_2!^M$ are names of different seriate sets of lines.

A set as simply an aggregation of objects all of the same the dimension is analogously indicated as such by the use of a subscript indicating the dimension of those objects. Thus $*_0S$ is a collection of points which has the Dimension 0, as distinct from $S_1!$ which indicates a *seriate set* of points which has the Dimension 1.

DEFINITION 2.1

Given any set of fundamental lines then the set of all (and only) the points within each one of those lines, is termed the set of points SUBSUMED by the given set of fundamental lines; and any point, or set of such points, within that subsumed set is said to be WITHIN the given set of fundamental lines.

DEFINITION 2.2

A fundamental line within the set of points subsumed by a given seriate set of fundamental lines $S_2!$, is a SERIATING line with respect to that set, if and only if, no two points within that given fundamental line are both within the same constituting line of $S_2!$.

CONVENTIONS 2.1

The name of a seriating line takes the form exemplified by $\underline{A} - \underline{B}(x)$. In any proof (or other context) where only one seriate set of fundamental lines is named, this syntax is taken to indicate that that seriating line has that characteristic with respect to that set.

DEFINITION 2.3

If, and only if, within a given seriate set, $S_N!$ a given object is an "E" object of each of two subsidiary seriate sets, $S_N!^A$ and $S_N!^B$, each within $S_N!$, and $S_N!^A$ and $S_N!^B$ have no other object in common, then the given object is BETWEEN the other two "E" objects of $S_N!^A$ and $S_N!^B$, within the given seriate set.

This definition simply generalises to all seriate sets the definition of between for points in a line given in In. 1.2. The syntax remains the same, except that a colon is introduced before the brackets containing the variable referencing the objects of the underlying set, where necessary to avoid ambiguity. E.g. for three lines in a seriate set of lines reference by 'x', $\underline{AB}(a) / \underline{CD}(b) / \underline{EF}(c) : (x)$

CONVENTION 2.2

The name of a seriate set of fundamental lines takes the form $S_2!(x, \underline{S_1!}^1, \underline{S_1!}^2)$, where 'x' references the fundamental lines constituting the given seriate set and $\underline{S_1!}^1$ and $\underline{S_1!}^2$ reference the "E" objects of that set.

THEOREM 2.1

In a seriate set of fundamental lines no point of any line in the set is an element of any other line in the set.

Definition 0.2(a)

THEOREM 2.2

Seriate sets of fundamental lines are self-similar both (a) externally and (b) internally in an exactly analogous way to fundamental lines.

Corollary: The between relationships which hold for points in a fundamental line hold for the constituting lines of a seriate set of fundamental lines.

Proofs in Th. 1.1 and 1.2 are unaffected if lines replace points in them, and the same follows for the theorems determining between relationships, given Th. 2.1.

DEFINITION 2.4

A given seriate set of fundamental lines is said to be UNFIXED if it is not part of any array within which there exists any object which has within it two points within different constituting line of that given set.

The notion of an unfixed seriate set allows us to consider the elements of that set purely in terms of the relationships that arise from their membership of that set, free from considerations as to what relationships might be asserted to exist between them otherwise.

THEOREM 2.3

Given an unfixed seriate set of fundamental lines together with any two points on two different constituting lines of that set, then the existence of a seriating line with those points as end-points, within the set of points subsumed by the given seriate set, can be asserted.

1. $S_2!(x)$
2. $[T \mapsto a] \iff \{[\underline{A}^M \underline{B}^M(x^M) \mapsto x] \& [T \mapsto x^M]\}$
3. $[\underline{A}^P \underline{B}^P(x^P) \& \underline{A}^Q \underline{B}^Q(x^Q) \mapsto x]$
4. $[P \mapsto x^P] \& [Q \mapsto x^Q]$
5. $S_2!^H(x^1, \underline{S}_1!^P, \underline{S}_1!^Q)$
 $\& [\underline{S}_1!^P = \underline{A}^P \underline{B}^P(x^P)] \& [\underline{S}_1!^Q = \underline{A}^Q \underline{B}^Q(x^Q)]$
 $\& [x^1 \mapsto x]$ 1,3,Th.2.2
6. $[J \mapsto g]$
7. $[J \mapsto x^N] \& [\underline{A}^N \underline{B}^N(x^N) \mapsto x^1]$ 6
8. $\neg \{[K \mapsto (g \& x^N)] \& \neg [J = K]\}$ 6,7
9. $[\underline{A}^N \underline{B}^N(x^N) \mapsto x^1] \implies \{J \& [J \mapsto (x^N \& g)]\}$

10. $[P \ \& \ Q] \mapsto g$
11. $*_0S(g) = \underline{PQ}(g)$ 6/10,5,D.0.3
12. $\underline{P-Q}(g) \ \& \ [g \mapsto a]$ 11,9,D.2.2

L. 2: Here 'a' is defined as a variable referencing all the points subsumed by the given seriate set. L. 6/10: Here 'g' is defined as a variable referencing only, one and only one point within every fundamental line within the seriate set $S!^H$, and including 'P' and 'Q'. L. 11: The set of points reference by 'g' has a one-to-one relationship with the seriate set of fundamental lines of $S!^H$, therefore those points cannot be distinguished from those of a seriate set hence that set is identical to a seriate set of points by D.0.3.

THEOREM 2.4

Given any unfixed seriate set of fundamental lines then the existence of a ring can be asserted constituted out of all and only the end-points of all the constituting fundamental lines of the given set together with all the points within the "E" objects of the set.

1. $S_2!(x, \underline{S_1!^1}, \underline{S_1!^2})$
 $\& [S_1!^1 = \underline{A^1B^1}(x^1)] \ \& [S_1!^2 = \underline{A^2B^2}(x^2)]$
2. $*_0S(f) \ \& \ *_0S(g) \ \& \ \neg \{P \ \& \ [P \mapsto (f \ \& \ g)]\}$
3. $\{[P = (A^N \wedge B^N)] \ \& \ [\underline{A^NB^N}(x^N) \mapsto x]\}$
 $\iff [P \mapsto (f \wedge g)]$
4. $[(A^1 \ \& \ A^2) \mapsto f] \ \& \ [(B^1 \ \& \ B^2) \mapsto g]$
5. $\underline{A^1A^2}(f) \ \& \ \underline{B^1B^2}(g)$ 3/4,Th.2.3,It,D.0.1
6. $A^1/A^2/B^2/B^1/A^1(r) \ \& \ \{r \mapsto [f + x^2 + g + x^1]\}$ 1,5,2,Th.1.1,ln.1.3

L. 2/3: Here the two variables 'f' and 'g' are defined as such that all, and only, the end-points of all the fundamental lines constituting the given seriate set are referenced by one or other of them but not by both. L. 4/5: The end-points of the fundamental lines constituting the "E" objects of the given seriate set are here arbitrarily assigned to 'f' and 'g' and then form the end-points of two fundamental lines referenced by those variables following the *rationale* of Th. 2.3. L. 6: Here one may be inclined to object that a ring cannot necessarily be so constituted because the lines referenced by 'f' and 'g' might intersect if the "E" objects of the seriate set were "the wrong way round." No such problem can exist however because the set is *unfixed* and hence there is nothing that can define any relationship between the members of the seriate set except membership of that set, so D. 0.1 governs: The situation must not be thought of as being one in which the lines of the given seriate set are "already" fixed on a page.

DEFINITION 2.5

The set of points constituting the ring as given in Th. 2.4 above is termed the BOUNDARY ((set of) points) of the set of points subsumed by the given seriate set.

DEFINITION 2.6

A FIXED seriate set of fundamental lines is identical to a seriate set of fundamental lines whose boundary set, with a defined membership, has been asserted to exist, and which is constituted out of the same objects as constitute the fixed set.

The notion of a fixed set is given in this indirect way in order to eliminate the need for formal specification of the boundary set and its membership, every time the existence of such a set is asserted. An unfixed and a fixed set are not directly converse: for example, if, relative to a given unfixed set, we first assert the existence of a fundamental line within that set constituted wholly out of the end-points of the constituting lines of the unfixed set, except in that one of its end-points is a non end-point of one of the constituting lines of the given seriate set, the unfixed set does not turn into a fixed set as here defined because then the ring so constituted as in Th. 2.4 cannot be asserted to exist within the resultant array.

THEOREM 2.5

Given a fixed seriate set of fundamental lines together with any two points on two different constituting lines of that set, then the existence of a seriating line with those points as end-points, within the set of points subsumed by the given seriate set, can be asserted.

This theorem is the exact repetition of Th. 2.3 except in so far as the given set is fixed rather than unfixed, and holds in the same way as the proof given there, and by D.0.1. This identical result which holds for both unfixed and fixed sets as defined does not in fact hold in the case of the somewhat unusual array described in the commentary to D. 2.6 above. By considering only unfixed and fixed sets, as they have been defined, we eliminate from direct consideration that rather anomalous set up.

THEOREM 2.6

Given any two non intersecting fundamental lines within the boundary set of a given seriate set of fundamental lines, then the existence of another seriate set of fundamental lines can be asserted, which set subsumes the same set of points as is subsumed by the given set and has the same boundary set as the first, and whose "E" objects are the two given fundamental lines.

1. $S_2!(x, \underline{S}_1!^1, \underline{S}_1!^2)$
 $\& [\underline{S}_1!^1 = \underline{A}^1 \underline{B}^1(x^1)] \& [\underline{S}_1!^2 = \underline{A}^2 \underline{B}^2(x^2)]$
2. $[P \mapsto a] \iff \{[\underline{A}^N \underline{B}^N(x^N) \mapsto x] \& [P \mapsto x^N]\}$
3. $\{[P = (A^N \wedge B^N)] \& [\underline{A}^N \underline{B}^N(x^N) \mapsto x]\}$
 $\iff [P \mapsto (f \wedge g)]$
4. $\neg \{P \& [P \mapsto (f \& g)]\}$
5. $\underline{A}^1 \underline{A}^2(f) \& \underline{B}^1 \underline{B}^2(g)$
6. $b \longleftrightarrow [f + x^2 + g + x^1]$
7. $\underline{J}^1 \underline{K}^1(y^1) \& \underline{J}^2 \underline{K}^2(y^2) \& [y^1 \mapsto f] \& [y^2 \mapsto g]$
 $\& A^1/J^1/K^1(f) \& B^1/J^2/K^2(g)$
8. $\underline{J}^1 \underline{J}^2(h) \& [h \mapsto b] \& [x^1 \mapsto h]$
 $\& \underline{K}^1 \underline{K}^2(w) \& [w \mapsto b] \& [x^2 \mapsto w]$

9. $[T^1 \mapsto a] \& \neg [T^1 \mapsto (y^1 \wedge y^2)]$
10. $\underline{J^3-K^3}(y^3, T^1) \& [y^3 \mapsto a] \& [J^3 \mapsto h] \& [K^3 \mapsto w]$
 $\& \{ \neg [P = (J^3 \wedge K^3)] \implies \neg [P \mapsto (y^3 \& b)] \}$
 $\& \neg \{ P \& [P \mapsto y^3] \& [P \mapsto (y^1 \wedge y^2)] \}$ 9,Th.2.5,D.0.1
11. $T^2 \& [T^2 \mapsto a] \& \neg [T^2 \mapsto (y^1 \wedge y^2 \wedge y^3)]$
12. $\underline{J^4-K^4}(y^4, T^2) \& [y^4 \mapsto a] \& [J^4 \mapsto h] \& [K^4 \mapsto w]$
 $\& \{ \neg [P = (J^4 \wedge K^4)] \implies \neg [P \mapsto (y^4 \& b)] \}$
 $\& \neg \{ P \& [P \mapsto y^4] \& [P \mapsto (y^1 \wedge y^2 \wedge y^3)] \}$ 11,Th.2.5,D.0.1
13. $*_1 S(y)$
14. $\{[(\underline{J^N K^N}(y^N) \& \underline{J^M K^M}(y^M)) \mapsto y] \& \neg [y^M \longleftrightarrow y^N]\}$
 $\implies \neg \{ P \& [P \mapsto (y^N \& y^M)] \}$
15. $\{P \& [P \mapsto a]\}$
 $\iff \{[\underline{J^N K^N}(y^N) \mapsto y] \& [P \mapsto y^N]\}$
16. $[P \mapsto s] \iff \{[\underline{J^N K^N}(y^N) \mapsto y] \& [P \mapsto y^N]\}$
17. $S_2!(y, \underline{S_1}!^3, \underline{S_1}!^4)]$
 $\& [S_1!^3 = \underline{J^1 K^1}(y^1)] \& [S_1!^4 = \underline{J^2 K^2}(y^2)]$
 $\& [s \longleftrightarrow a]$ 13/16,9/12,lt.T(a),D.0.3
18. QED 17,lt.

Note that in the statement of the theorem, because the boundary set is given we do not need to state the fixed condition. NON INTERSECTING lines have no point in common. Intersecting lines have at least one point in common. L. 1/8 describe the set-up. L. 10 & 12: The seriating lines formed as in Th. 2.5 by taking one point from each of the constituting lines of $S!(x)$ between the two constituting lines of that set within which their end-points lie. L. 17: By iterating for every possible 'T' in 'a' as in L. 9/10,11/12 then the existence of a set of non intersecting lines $*_1(y)$ can be asserted which set is not distinguishable from a seriate set, hence L. 17. L. 18: reference again to L. 9/12 shows the boundary set of the set so formed to be identical to that of the given set. If the given pair of fundamental lines are not situated as in L.7 a small number of iterations of L. 17 on judiciously selected intermediate pairs of lines achieves the required result.

THEOREM 2.7

Any two seriating line within a fixed seriate set of fundamental lines with all their end-points within two of the constituting lines of that set and being non intersecting or, at most having solely in common one end-point, constitute the boundary set of seriate set of lines.

This follows from the iteration of Th. 2.2(b) and the use of the method of Th. 2.6. Where the seriating lines have an end-point in common, that common point should not be within the fundamental lines selected as the "E" objects of the required set.

THEOREM 2.8

Given any seriate set of fundamental lines then any fundamental line within the set intersects every constituting line of that given set which is between any two constituting lines of the set within which lie points of the given line.

1. $S_2!(x, \underline{S}_1!^1, \underline{S}_1!^2)$
2. $[J \mapsto a] \iff \{[\underline{A}^N \underline{B}^N(x^N) \mapsto x] \& [J \mapsto x^N]\}$
3. $[\underline{S}_1!^1 = \underline{A}^1 \underline{B}^1(x^1)] \& [\underline{S}_1!^2 = \underline{A}^2 \underline{B}^2(x^2)]$
4. $\underline{PQ}(p) \& [p \mapsto a]$
 $\& [P^1 \mapsto (p \& x^P)] \& [Q^1 \mapsto (p \& x^Q)]$
 $\& \{[\underline{A}^Q \underline{B}^Q(x^Q) \& \underline{A}^P \underline{B}^P(x^P)] \mapsto x\}$
5. $\underline{A}^P \underline{B}^P(x^P) / \underline{A}^N \underline{B}^N(x^N) / \underline{A}^Q \underline{B}^Q(x^Q) : (x)$
6. $[J \mapsto f] \iff$
 $\{[\underline{A}^1 \underline{B}^1(x^1) / \underline{A}^M \underline{B}^M(x^M) / \underline{A}^N \underline{B}^N(x^N) : (x)$
 $\& (J \mapsto x^M)] \wedge [J \mapsto x^1]\}$
7. $[J \mapsto g] \iff$
 $\{[\underline{A}^N \underline{B}^N(x^N) / \underline{A}^M \underline{B}^M(x^M) / \underline{A}^2 \underline{B}^2(x^2) : (x)$
 $\& (J \mapsto x^M)] \wedge [J \mapsto x^2]\}$
8. $\neg \{J \& [J \mapsto (f \& g)]\}$ 6,7,Th.2.2Cr,1.3
9. $\neg [T \& [T \mapsto (x^N \& p)]]$
10. $[K \mapsto p] \implies [K \mapsto (f \wedge g)]$ 9,6,7,1,4
11. $Z \& [Z \mapsto p] \& \neg [Z \mapsto (f \& g)]$ 8,10,D.0.2
12. $T \& [T \mapsto (x^N \& p)]$ 10,11

Corollary: Any fundamental line within a seriate set of fundamental lines with end-points one to each of the "E" objects of that set, intersects every constituting line of that set.

L. 1/4 describe the set-up and L. 5 defines a constituting line of the seriate set between the two constituting lines of that set within which lie the two given points within the given fundamental line. L. 6,7 define the two sets of all the points "on either side" of that "between" line, which, by Th. 2.2,Cr. and Th. 1.3 applied to every pair of one from each have no point in common (L. 8). L. 9 assumes that the "between" line does not intersect the given fundamental line. Thus all its points must lie within the sets of points "to either side" of the "between" line (L. 10). But there must then be a point between every pair of points one from 'f' and one from 'g' within the given line, ergo there must be a point that line not referenced by either, in contradiction of this last (L. 11). So there must be a point of the fundamental line within the "between" line (L. 12).

THEOREM 2.9

Given any fundamental line within a fixed seriate set of lines then the existence of a *minimal* set of *seriatly catenated* fundamental lines can be asserted, which set of lines subsumes the same set of points as are subsumed by the given lines, and whose members are each either wholly within a constituting line of the seriate set of lines, or are seriating lines with respect to that set.

1. $S_2!(x, \underline{S}_1!^1, \underline{S}_1!^2)$
2. $[T \mapsto a] \iff \{[\underline{A}^N \underline{B}^N(x^N) \mapsto x] \ \& \ [T \mapsto x^N]\}$
3. $\underline{PQ}(p) \ \& \ [p \mapsto a]$
4. $[J \mapsto x^J] \ \& \ [\underline{A}^J \underline{B}^J(x^J) \mapsto x]$
5. $J \mapsto [n^P \wedge n^Q]$
6. $\{[J \mapsto n^P] \implies [E = P]\} \ \& \ \{[J \mapsto n^Q] \implies [E = Q]\}$
7. $F \ \& \ \neg \{F^1 \ \& \ F/F^1/J(p) \ \& \ \neg [F^1 \mapsto x^J]\} \ \& \ E/F/J(p) \quad 4/6$
8. $[J \mapsto n^B] \implies [J \mapsto (n^P \ \& \ n^Q)]$
9. $J \mapsto s^B$
10. $F \ \& \ [F \mapsto S_1!(x^F)] \ \& \ G \ \& \ [G \mapsto S_1!(x^G)]$
 $\quad \& \ S_1!(x^F) / S_1!(x^J) / S_1!(x^G) : (x)$
 $\quad \& \ \neg \{[F^1 \ \& \ F/F^1/J(p) \ \& \ (F^1 \mapsto x^J)]\}$
 $\quad \& \ \neg \{[G^1 \ \& \ G/G^1/J(p) \ \& \ (G^1 \mapsto x^J)]\} \quad 9,4$
11. $[T \mapsto k] \implies \{[T \mapsto p] \ \& \ \neg [T \mapsto (n^B \wedge s^B)]\}$
12. $[\underline{T}^N \underline{T}^M(p^N) \mapsto c] \iff$
 $\quad \{[(T^N \ \& \ T^M) \mapsto k] \ \& \ [p^N \mapsto p]$
 $\quad \& \ \neg [(T^Z \mapsto k) \ \& \ T^M/T^Z/T^N(p)]\}$
13. $\underline{Z}^N \underline{Z}^M(p^Z) \mapsto c$
14. $\{\underline{Z}^N \underline{Z}^M(p^Z) = \underline{Z}^N - \underline{Z}^M(p^Z)\}$
 $\quad \wedge \ \{[p^Z \mapsto x^N] \ \& \ [\underline{A}^N \underline{B}^N(x^N) \mapsto x]\} \quad 13,12$
15. $\text{QED } 13/14,12$

A SERIATELY CATENATED set of fundamental lines conforms to *S in D. 0.2(b)ii/iv, and a minimal such set is the set with the least number of members which conforms to some other stated condition(s). L. 5/8: n^B points are defined as points at which the given line which "continues towards" both 'P' and 'Q' within a constituting line of the underlying seriate set. L. 9/10 define s^B points as where the given line "traverses" the underlying seriate set "in opposite directions." L. 11: 'k' then references all the points of the given line not referenced by n^B or s^B , that is are either its end-points or are points where it "varies its traverse" with respect to the underlying set. L. 12: 'c' here is defined as referencing all the "non overlapping" fundamental lines within

the given line into which the 'k' point "divide it." These lines must then be constituted by "l" points which are either all n^B points and hence are within a constituting line of the given seriate set, or by s^B points and hence are seriating lines with respect to it, because if not, any such line would have "a kink" in it contrary to L. 12.

INTERPRETATION 2.2: FUNDAMENTAL AREAS

A set of points constitutes a FUNDAMENTAL AREA, if and only if, it is identical one-to-one to the set of points subsumed by a fixed seriate set of fundamental lines. The name of an area is exemplified by $A!(a)$ where 'a' references all the points subsumed by that seriate set of fundamental lines. The addition of an underlined variable within the round brackets, as for example by 'b' in $A!(a, \underline{b})$, references the boundary (set of) points of the area identical to the boundary set relative to which the seriate set of lines is fixed.

THEOREM 2.10

Given any two fundamental areas, having in common solely a set of boundary points which constitutes a single fundamental line, then the existence of a fundamental area can be asserted which subsumes all and only the points subsumed by the two given fundamental areas, and whose boundary set is constituted out of the sum of the boundary sets of the two given areas less the non end-points of the fundamental line common to both.

In accord with In. 2.2 and Th. 2.6 the existence can be asserted of two seriate sets of fundamental lines, each equivalent to one of the two given areas, which seriate sets both have as an "E" object the fundamental line common to the two given areas. By Th. 2.2(a) these sets conjoin to form a seriate set of fundamental lines which is equivalent to an area subsuming all the points subsumed by the original two areas, and with a border as given.

THEOREM 2.11

Given any fundamental line within a fundamental area with solely its end-points within the boundary of the given area, then the existence of two fundamental areas within the given fundamental area can be asserted, which areas together subsume the same set of points as is subsumed by the given area and which have no points in common except the points within the given fundamental line.

1. $A!(a^1, \underline{b}^1)$
2. $\underline{JK}(q) \ \& \ [q \mapsto a^1] \ \& \ [(J \ \& \ K) \mapsto b^1]$
 $\ \& \ \{ \neg [P = (J \wedge K)] \implies \neg [P \mapsto (q \ \& \ b^1)] \}$
3. $S_2!(x, \underline{S}_1!^1, \underline{S}_1!^2)$
 $\ \& \ [S_1!^1 = \underline{A^1B^1}(x^1)] \ \& \ [S_1!^2 = \underline{A^2B^2}(x^2)]$
 $\ \& \ [(x^1 \ \& \ x^2) \mapsto b^1] \ \& \ [J \mapsto x^1] \ \& \ [K \mapsto x^2]$
4. $[P \mapsto a^1] \iff \{[\underline{A^N B^N}(x^N) \mapsto x] \ \& \ [P \mapsto x^N]\}$ 1,3,In.2.2,Th.2.6
5. $[P \mapsto f] \iff$
 $\ \{[P = (J^1 \wedge K^1)] \ \& \ \underline{J^1 - K^1}(q^1) \ \& \ [q^1 \mapsto q] \ \&$

- $\neg [\underline{J^-} K^2(q^2) \& (q^1 \mapsto q^2) \& \neg (q^1 \leftrightarrow q^2)]\}$
6. $[Q \mapsto g] \iff$
 $\{P \& [P \mapsto ({}^f \& x^N)] \& [\underline{A^N B^N}(x^N) \mapsto x]$
 $\& [Q \mapsto x^N] \& [Q \mapsto (q \wedge b^1)]\}$
 7. $[\underline{J^1 K^1}(q^1) \mapsto m] \iff$
 $\{[\underline{J^1-} K^1(q^1) \& [q^1 \mapsto (q \wedge b^1)] \& [(J^1 \& K^1) \mapsto g]$
 $\& \neg [J^1/W/K^1(q^1) \& (W \mapsto g)]\}$
 8. $[\underline{N^1 N^2}(n^1) \mapsto n] \implies$
 $\{[\underline{A^N B^N}(x^N) \mapsto x] \& [n^1 \mapsto x^N]\}$
 9. $A!(a^S, \underline{r^S}) \mapsto s$
 10. $[\underline{M^1 N^1}(m^1) \& \underline{M^2 N^2}(m^2)] \mapsto m$
 11. $[\underline{M^1 M^2}(n^1) \mapsto n] \& \{[\underline{N^1 N^2}(n^2) \mapsto n] \wedge [N^1 = N^2]\}$
 12. $\neg \{[\underline{M^3 N^3}(m^3) \mapsto m] \& [M^3 \mapsto n^1] \& [N^3 \mapsto n^2]$
 $\& M^1/M^3/M^2(n^1)\}$
 13. $\{\neg [N^1 = N^2] \implies [r^S \leftrightarrow [m^1 + n^2 + m^2 + n^1]$
 $\& \{[N^1 = N^2] \implies [r^S \leftrightarrow [m^1 + m^2 + n^1]]\}$
 $\& \{M^1/N^1/N^2/M^2/M^1(r^S) \wedge M^1/N^1/M^2/M^1(r^S)\}$
 $\& [a^S \mapsto a^1]\}$ 9,Th.1.1,Th.2.7
 14. $[\underline{M^1 N^1}(m^1) \mapsto m] \& [m^1 \mapsto q]$
 15. $\{[A!(a^N, \underline{r^N}) \& A!(a^M, \underline{r^M})] \mapsto s\}$
 $\& [m^1 \mapsto (r^N \& r^M)] \& \neg [r^N \leftrightarrow r^M]$
 $\& [A!(a^N, \underline{r^N}) \mapsto c] \& [A!(a^M, \underline{r^M}) \mapsto d]$ 14
 16. $\{[A!(a^N, \underline{r^N}) \& A!(a^M, \underline{r^M})] \mapsto s\} \& \neg [a^M \leftrightarrow a^N]$
 $\& [\underline{M^1 M^2}(n^M) \mapsto n] \& [n^M \mapsto (r^N \& r^M)]$
 17. $\{[A!(a^N, \underline{r^N}) \mapsto c] \implies [A!(a^M, \underline{r^M}) \mapsto c]\}$
 $\& \{[A!(a^N, \underline{r^N}) \mapsto d] \implies [A!(a^M, \underline{r^M}) \mapsto d]\}$ 16
 18. $[P \mapsto a^2] \iff \{[P \mapsto a^N] \& [A!(a^N, \underline{r^N}) \mapsto c]\}$
 19. $[P \mapsto a^3] \iff \{[P \mapsto a^N] \& [A!(a^N, \underline{r^N}) \mapsto d]\}$
 20. $A!(a^2, \underline{b^2}) \& A!(a^3, \underline{b^3})$

$$\begin{aligned}
& \& \{ [P \mapsto (b^2 \wedge b^3)] \implies [P \mapsto (q \wedge b^1)] \} \\
& \& \neg \{ P \& \neg [P \mapsto q] \& [P \mapsto (a^2 \& a^3)] \} \\
& \& [a^1 \longleftrightarrow (a^2 + a^3)] \qquad \text{Th.2.9,Th. 2.10,lt.}
\end{aligned}$$

This theorem together with the last show how fundamental areas are externally and internally self-similar. L. 5: 'f' references the end-points of all the seriating lines of a minimal seriate catenation equivalent to the given line. L. 6: 'g' references every intersection point of every constituting line of the given seriate set having an 'f' point within it, with the given line, or the with boundary of the given area. L. 7 defines all the seriating lines within the given line or the boundary which have 'g' points as end-points and no similar point between them. L. 9/13 define as 's' areas all the areas within the given area defined by any two 'm' lines with end-points on the same two constituting lines of the underlying seriate set, and with no such line between them. L. 14/15: Each of the two 's' areas "lying on opposite sides" of every seriating section of the given line are then assigned each to one of two mutually exclusive classes, those areas "lying side-by-side" being assigned to the same class (L. 16/17). All the areas in each one of these classes are then added together in accord with Th. 2.10 to form two areas with Th. 2.9 ensuring they are together equivalent to the original area. If the consistency of the assignment of areas to 'c' and 'd' is doubted, application of Th. 2.8 shows the agreement between L. 15 & L. 17.

THEOREM 2.12

Given a ring within a fundamental area then the existence can be asserted of one and only one fundamental area within the given fundamental area with that ring as its boundary (set).

The existence of the area with the given ring as boundary follows from a maximum of two, judicious partitionings of the given fundamental area in accord with Th. 2.11. If another area with the same boundary set could exist within the same given area then the assignment of points by Th. 2.11 would be controverted.

THEOREM 2.13

Given any pair of fundamental areas having in common only boundary points, together with a fundamental line wholly within those areas, and having a non boundary point of each within it, then a point within that fundamental line must also be a common boundary point of those areas.

If the two areas have merely one point in common then the given fundamental line must have that point within it or it can not be such by D. 0.2. Otherwise, iterated judicious distinction of seriate sets equivalent to the given areas and application of Th. 2.8 governs.

THEOREM 2.14

If three fundamental lines within a given fundamental area have solely in common a pair of points which are the end-points of all of them, then one and only one pair of those three fundamental lines forms the boundary set of a fundamental area whose existence within the given fundamental area can be asserted, which area subsumes all the points within the other two fundamental areas whose existence can be asserted within the given area, defined by boundary sets constituted out of the other two pairs of lines from the original three.

1. $A!(a)$
2. $\underline{PQ}(x) \& \underline{PQ}(y) \& \underline{PQ}(z) \& [(x \& y \& z) \mapsto a]$

3. $\neg [T = (P \wedge Q)]$
4. $\neg [T \mapsto [(x \& y) \wedge (x \& z) \wedge (y \& z)]]$ 3
5. $A!(a^1, \underline{b}^1) \& [a^1 \mapsto a] \& [b^1 \leftrightarrow (x + y)]$ 1,2,Th.2.12
6. $A!(a^2, \underline{b}^2) \& [a^2 \mapsto a] \& [b^2 \leftrightarrow (x + z)]$ 1,2,Th.2.12
7. $A!(a^3, \underline{b}^3) \& [a^3 \mapsto a] \& [b^3 \leftrightarrow (z + y)]$ 1,2,Th.2.12
8. $[X \mapsto x] \& [Y \mapsto y] \& [Z \mapsto z]$
 $\& \neg [(X \wedge Y \wedge Z) = (P \wedge Q)]$
9. $[Z \mapsto a^1]$
10. $[(a^2 \& a^3) \mapsto a^1] \& [Z \mapsto (a^1 \& a^2 \& a^3)]$ 9, Th.2.13
11. $\neg [X \mapsto a^3] \& \neg [Y \mapsto a^2]$ 9,Th.2.12,2.11
12. $\neg [Z \mapsto a^1] \implies \{[X \mapsto a^3] \wedge [Y \mapsto a^2]\}$ Th.2.12,2.11
13. QED 9/11,11,Sym.

L. 10: As 'Z' a non end-point of the 'z' line is within a^1 then every point of that line must be within a^1 or else Th. 2.13 is controverted by L. 3/4 and similarly every point of a^2 must be within a^1 . L. 11: Th. 2.11 and Th. 2.12 require that given L. 9/10 a unique layout arises with 'X' and 'Y' in different subsidiary areas. L. 12: If 'Z' is not in a^1 then, if 'X' is not in a^3 , then add a^1 to a^3 to form a^2 and 'Y' must be within it, and this covers all possibilities.

THEOREM 2.15

No three fundamental areas all within a given fundamental area, which have solely boundary points in common, can all have the same one point in common, which point is a non end-point of a fundamental line constituted out of boundary points of two of them.

Assume such a point exists. Then consider a ring within the conjunction of the two areas which have the fundamental line in common, with the postulated point as a non boundary point of the area within that ring. If a third area, not within the other two, has that point within it then by Th. 2.5 the existence of a fundamental line can be asserted having within it the postulated point and also a point not within that ring, which line does not intersect the ring, in contravention of Th. 2.13.

A Five Country Map

Five fundamental areas, each within a given fundamental area, such that they have no points in common except boundary points, cannot have each with every other a common set of points which constitutes a fundamental line.

1. $A!(a)$
2. $\underline{P^N P^M}(j) \mapsto t$
3. $A!(a^N, \underline{b}^N) \& A!(a^M, \underline{b}^M)$

- $$\begin{aligned}
& \& \neg[a^M \longleftrightarrow a^N] \& [(a^N \& a^M) \longmapsto a] \\
& \& \{\neg[Z \longmapsto (b^N \& b^M)] \implies \neg[Z \longmapsto (a^N \& a^M)]\} \\
& \& \underline{FG}(k) \& [k \longmapsto (b^N \& b^M)] \\
4. \quad & [P^N \longmapsto a^N] \& \neg[P^N \longmapsto b^N] \\
& \& [P^M \longmapsto a^M] \& \neg[P^M \longmapsto b^M] \\
& \& [j \longmapsto (a^N + a^M)] \\
& \& [T \longmapsto (j \& k)] \& \neg[T = (F \wedge G)] \\
& \& \{\neg[T^1 = T] \implies \neg[T^1 \longmapsto (j \& k)]\} \\
5. \quad & \{[\underline{P^N P^1}(j^1) \& \underline{P^N P^2}(j^2)] \longmapsto t\} \& \neg[j^1 \longleftrightarrow j^2] \\
6. \quad & \neg[T = P^N] \implies \neg[T \longmapsto (j^1 \& j^2)] \quad 5,D.0.1 \\
7. \quad & \{ \underline{P^A P^C}(j^A C) \& \underline{P^C P^B}(j^C B) \} \\
& \& \underline{P^A P^D}(j^A D) \\
& \& \underline{P^D P^B}(j^D B) \\
& \& \underline{P^A P^E}(j^A E) \\
& \& \underline{P^E P^B}(j^E B) \\
& \& \underline{P^A P^B}(j^A B) \} \longmapsto t \\
8. \quad & \underline{P^A P^B}(w^C) \& \underline{P^A P^B}(w^D) \& \underline{P^A P^B}(w^E) \\
& \& [w^C \longleftrightarrow (j^A C + j^C B)] \\
& \& [w^D \longleftrightarrow (j^A D + j^D B)] \\
& \& [w^E \longleftrightarrow (j^A E + j^E B)] \\
& \& [w \longleftrightarrow j^A B] \quad 7,Th.1.1 \\
9. \quad & \dots /P^A//P^B/\dots(h^1) \& [h^1 \longleftrightarrow (w^C + w^E)] \\
& \& A!(f^1, \underline{h}^1) \& [f^1 \longmapsto a] \\
& \& [P^D \longmapsto a^1] \quad 5/6,8,Th.2.14,Rn. \\
10. \quad & \dots /P^A//P^B/\dots(h^2) \& [h^2 \longleftrightarrow (w + w^C)] \\
& \& A!(f^2, \underline{h}^2) \& [f^2 \longmapsto a] \\
& \& [P^E \longmapsto f^2] \quad 5/6,8,Th.2.14,Rn. \\
11. \quad & \neg\{\underline{P^C P^E} \& [\underline{P^C P^E} \longmapsto t]\} \quad 10,Th.2.13,2.15 \\
12. \quad & \dots /P^A//P^B/\dots(h^3) \& [h^3 \longleftrightarrow (w + w^E)]
\end{aligned}$$

$$\& A!(f^3, h^3) \& [f^3 \mapsto a^1]$$

$$\& [P^D \mapsto f^3]$$

5/6,8,Th.2.14,Rn.

$$13. \quad \neg \{ \underline{P^C P^D} \& [\underline{P^C P^D} \mapsto t] \}$$

12,Th.2.13,2.1

14.

QED 11,13,Sym.

L. 2/4: 't' is set up as referencing any fundamental line such as must exist when (L. 3) two different areas are within a given area, which areas have solely boundary points in common, and have a fundamental line in common; which line referenced by 't' (L. 4) has as end-points a non boundary point of each of the two different areas, is wholly within them and intersects their common boundary line, not in its end-points, in one and only one point (this last simply to aid metal clarity). L. 5/6: Any two 't' lines with a common end-point are stipulated to have no other point in common. L. 7: A number of 't' lines are then set out relative to five areas - merely alluded to by superscripts A,B,C,D,E, because they do not require formal identification, in order to aid comprehension. These 't' lines "connect" two base areas 'A' and 'B' to each of the other three. L. 8/9: By suitable additions a symmetrical triplet of lines, each line connecting the two base areas through one of the others, is formed, on the pattern of Th. 2.14. By renaming, the situation is as given. L. 10: The 't' line "connecting" the two base areas directly is then assumed to be not within a^1 . But then P^E is within, and P^C is not within, a ring consisting wholly of 'D', 'A' and 'B' points and hence by Th. 2.13 and 2.15 a 't' line "connecting" 'E' and 'C' cannot exist (L. 11). Similarly if the 'w' line is within a^1 , the same problem arises between 'D', and 'E' or 'C' (L. 12/13). By symmetry all situations have now been examined so 't' lines cannot exist each "connecting" one of every pair out of five areas, which would have to be the case if they each had with every other, a boundary line section in common.

NOTES

The abbreviations used to refer to different subsections in the paper are as follows: D. = Definition; Cn. = Convention; In. = Interpretation; Th. = Theorem; Cr. = Corollary. In the third columns of proofs It. stands for iteration; Rn. for rename; Sim. for similarly or similarity; Sym. for symmetry. In the commentary following proofs L. = Line.

Cn. 1.1: The relationship between the objects referenced by a variable and that variable may be thought of as roughly similar to the relationship between a mathematical variable and the values it takes. However no great weight should be placed on any such analogy. The usage of the calculus demonstrates the nature of the concept being employed.

The structure of proofs: Proofs are set out as numbered "lines" each consisting of a statement which takes a truth value (such "lines" may occupy more than one line on the page, but this is merely to facilitate the grouping of interlinked relationships, in order to aid comprehension.) The third column of a proof indicates something of its structure, typically by listing previous lines as one or more premises on which the current line depends, and often giving in addition the number of a relevant theorem which justifies that entailment, or other related information as alluded to in the abbreviations given above. Where a colon appears in this column it indicates one entailment embedded in another. However this third column is not set out in accord with any set of rigorous rules from which the exact structure of the proof can be extracted. The commentary which follows most proofs should be referred to, in order to establish the proof's structure. Typically omitted from the third column are the relationships indicated in the commentary to Th. 1.1; and generally where a previous line simply serves to define a variable or object, it is not listed.

Continuity: It may be asked what happens when you take an end-point, say, away from a fundamental line? What is left? Certainly the rules of calculus allow that given $\underline{AB}(x)$ one can assert the existence of $*S(y)$ and $y = [x - A]$. One is then inclined to shift to the interpretive level and say, well what sort of "object" does 'y' represent then? This however is an illegitimate move. 'y' cannot function as a variable referencing the points in a fundamental line because whatever end-point 'Z' one takes to accompany 'B' to form the end-points of that line, there always remains a point referenced by 'y' between 'A' and 'Z' and hence not in that fundamental line. The set $*S(y)$ then remains just that, a mere unstructured aggregation of points which cannot constitute a fundamental object of the calculus. This is but one of the ways continuity, which is ruptured by such a procedure, is expressed by the calculus.

Cn. 1.1: The syntax $[A \mapsto \neg x]$ is not allowed because it can all too easily lead to a drift into thinking of $\neg x$ as an *object of the calculus* where this is illegitimate. For example consider two fundamental areas, with no boundary points in common, the first having within it the second. If 'a' references the points of the "contained" area, to think of points "outside" the "contained" area as mapping onto those points of the "containing" area which are "not a", tends to attract one into thinking of "not a" as referencing the points of some kind of object or collection of objects of the calculus. But those points cannot constitute a well-formed array of the calculus analogously to the case for a line noted above. Two fundamental areas suitably conjoined and together having within them *the boundary points* of the "contained" area, are necessary, in order to constitute the outer annulus of the situation described. There thus seems to be two ways of constructing the same situation, either as an annulus or as one area within another, which two different descriptions "compete for" the same set of points, to form either the internal border of the annulus or the boundary of the contained area. This situation in fact appears to have a strong resonance with the well known *gestalt shift* between figure and ground: that is the alternate perceptions of the same visual field as constituting either an object lying on a surface, or an annular ring (against some background). Whether any such parallel is of any significance however remains to be seen.

Cn. 2.1: If in order to avoid confusion it is necessary to indicate the seriate set relative to which a line is a seriating line one can do this for example by the syntax $\underline{A-r-B}(x)$ where "r" is the variable referencing the constituting lines of the relevant seriate set.

APPENDIX

The purpose of this appendix is to set out more rigorous proofs of Th. 1.1 and 1.2 of the main text. In order to minimise repetition *syntactic variables* will be employed. *Syntactic variables* are variables which are defined in a

formal way in preliminary section, as referencing sets and/or objects all having the same, often complex, interlinking characteristics. In proofs these variables can then be directly alluded to in order to assign those same sets of characteristics directly to different objects without the need to list all those same characteristics every time. These variables are indicated by the inclusion of '%' as an element of their names, as in $s\%$ and $e^C\%(n)$.

DEFINITION SV1: Syntactic variables $s\%, e^E\%, e^C\%, p\%$

1. $[*S(n) \mapsto s\%] \implies \{ *S_1(m) \ \& \ [m \longleftrightarrow n] \}$
2. $\underline{J^N K^N}(j^N) \mapsto n$
3. $\{ [P \mapsto j^N] \ \& \ \neg [P = (J^N \wedge K^N)] \} \implies r \neg \{ [P \mapsto j^M] \ \& \ [\underline{J^M K^M}(j^M) \mapsto n] \ \& \ \neg [j^N \longleftrightarrow j^M] \}$ 1,2
4. $P \mapsto e^E\%:n$
5. $\{ [P = (J^M \wedge K^M)] \ \& \ [\underline{J^M K^M}(j^M) \mapsto n] \ \& \ \neg \{ [P \mapsto j^L] \ \& \ [\underline{J^L K^L}(j^L) \mapsto n] \ \& \ \neg [j^L \longleftrightarrow j^M] \} \}$ 1,4
6. $P \mapsto e^C\%:n$
7. $[P = (J^M \wedge K^M)] \ \& \ [P = (J^L \wedge K^L)] \ \& \ \{ [\underline{J^M K^M}(j^M) \ \& \ \underline{J^L K^L}(j^L)] \mapsto n \} \ \& \ \neg [j^L \longleftrightarrow j^M] \ \& \ \neg \{ [P \mapsto j^R] \ \& \ [\underline{J^R K^R}(j^R) \mapsto n] \ \& \ \neg [j^R \longleftrightarrow (j^M \wedge j^L)] \}$ 1,6
8. $\{ [P = (J^M \wedge K^M)] \ \& \ [\underline{J^M K^M}(j^M) \mapsto n] \} \implies [P \mapsto (p^E\%:n \wedge p^C\%:n)]$ 1
9. $[P \mapsto p\%:n] \iff \{ [\underline{J^M K^M}(j^M) \mapsto n] \ \& \ [P \mapsto j^M] \}$

Here the syntactic variable $s\%$ is defined as referencing a set of fundamental lines, which have no points in common except end-points, and whose each end-point is either, an element of one and only one other of the lines of the set, or of no other such line; and whose total set of subsumed points is referenced by $p\%:n$.

DEFINITION SV2: $w\%$

1. $*S(n) \mapsto w\%$
2. $*S(n) \mapsto s\%$ 1
3. $P^1 \ \& \ P^2 \ \& \ \{ [P^1 \ \& \ P^1] \mapsto e^E\%:n \} \ \& \ \neg [P^1 = P^2] \ \& \ \neg \{ P^3 \ \& \ [P^3 \mapsto e^E\%:n] \ \& \ \neg [P^3 = (P^1 \wedge P^2)] \}$ 1

$w\%$ references a set of fundamental lines which, in addition to being referenced by $s\%$, are such that two and only two of the end-points of those lines are referenced by $e^E\%$. As a result such a set conforms to the requirements of D. 0.2(b) clauses (ii)/(iv) for $*S$ there.

SV1/SV2 CONVERSE

If the entailments resulting from any set mapping onto either $s\%$ and $w\%$ are fulfilled by any set then that set maps onto those variables.

THEOREM F1: Given two fundamental lines $\underline{AB}$ and $\underline{BC}$ which have no point in common except 'B', then the existence of a fundamental line $\underline{AC}$, constituted out of all and only the sum of the points of $\underline{AB}$ and $\underline{BC}$, can be asserted.

1. $\underline{AB}(x) \ \& \ \underline{BC}(y)$
2. $\neg \{ [P \mapsto (x \ \& \ y)] \ \& \ \neg [P = B] \}$
3. $R \ \& \ [R \mapsto x] \ \& \ \neg [R = (A \wedge B)]$
4. $*S(n) \ \& \ [*S(n) \mapsto w\%] \ \& \ [R \mapsto e^C\%:n]$
 $\quad \quad \quad \& \ [(A \ \& \ B) \mapsto e^E\%:n] \ \& \ [x \longleftrightarrow p\%:n]$ 3,1,D.0.2,SV.1/2
5. $z \longleftrightarrow (x + y)$
6. $*S(m) \ \& \ [*S(m) \mapsto s\%] \ \& \ \{m \longleftrightarrow [n + \underline{BC}(y)]\}$
 $\quad \quad \quad \& \ [(B \ \& \ R) \mapsto e^C\%:m] \ \& \ [(A \ \& \ C) \mapsto e^E\%:m]$
 $\quad \quad \quad \& \ [z \longleftrightarrow p\%:m]$ 1,2,4,5)
7. $*S(m) \mapsto w\%$ 6,SV.1/2,Conv.
8. $*S(h) \ \& \ [*S(h) \mapsto w\%] \ \& \ \{h \longleftrightarrow [\underline{AB}(x) + \underline{BC}(y)]\}$
 $\quad \quad \quad \& \ [B \mapsto e^C\%:h] \ \& \ [(A \ \& \ C) \mapsto e^E\%:h]$
 $\quad \quad \quad \& \ [z \longleftrightarrow p\%:h]$ 1,2,SV.1/2,Conv.
9. $\underline{AC}(z) \ \& \ [z \longleftrightarrow (x + y)]$ 7,It(R),Sm,8,D.0.2


L. 9: The result of L. 7 occurs for every 'R' in 'x' and similarly for every non end-point of 'y'. L. 8 gives the same result for 'B'. Thus the requirement of a seriate set for every non end-point of 'z' is fulfilled. Here clause (v) of D. 0.2(b) is ignored. In fact this clause is not strictly necessary to the definition of a seriate set and is only included in D. 0.2 to allow the intuitive syntax in the introductory proofs of Th. 1.1 & 1.2 of the main text, to be directly employed. Clearly Th. F1 still goes through when that clause is ignored - any such ring as it precludes simply existing before and after. Then, by application of Th. F1 itself and the axiom of stability of D. 0.4 the requirement of that clause can be proved. Intuitively this may seem wrong, as if the truth of clause (v) must be proved before Th. F1 can be proven. However here as elsewhere it is the entailments of the calculus which must be attended to, not the interpretive situation where we may feel compelled to think that the intuitive syntax used in the main text must be justified before any such addition can occur.

THEOREM F2: Given a non end-point C within a fundamental line $\underline{AB}$ then the existence of two fundamental lines $\underline{AC}$ and $\underline{CB}$ within $\underline{AB}$ can be asserted, having C as their sole common point, and being constituted by all and only the same set of points as constitute $\underline{AB}$.


1. $\underline{AB}(x)$
2. $[C \mapsto x] \ \& \ \neg [C = (A \wedge B)]$
3. $*S(n) \ \& \ [*S(n) \mapsto w\%] \ \& \ [(A \ \& \ B) \mapsto e^E\%:n]$
 $\quad \quad \quad \& \ \{ [P \mapsto t] \iff [P \mapsto e^C\%:n] \} \ \& \ [C \mapsto t]$
 $\quad \quad \quad \& \ [p\%:n \longleftrightarrow x]$ 1,2,D.0.2,SV.1/2
4. $\{ [*S(j) \ \& \ *S(k)] \mapsto w\% \}$
 $\quad \quad \quad \& \ [(j + k) \longleftrightarrow n] \ \& \ \neg \{ S_1! \ \& \ [S_1! \mapsto (j \ \& \ k)] \}$
 $\quad \quad \quad \& \ [(A \ \& \ C) \mapsto e^E\%:j] \ \& \ [(C \ \& \ B) \mapsto e^E\%:k]$

- $$\begin{aligned}
& \& \{[P \mapsto t^1] \iff [P \mapsto e^C\%:j]\} \\
& \& \{[P \mapsto t^2] \iff [P \mapsto e^C\%:k]\} \\
& \& \{[t - C] \iff [t^1 + t^2]\} \\
& \& [(p\%:j + p\%:k) \iff x] \qquad \qquad \qquad 3,SV.1/2,Conv. \\
5. \quad & [p\%:j \iff y] \& [p\%:k \mapsto z] \\
6. \quad & \underline{AC}(y) \& \underline{CB}(z) \\
& \& [x \iff (y + z)] \\
& \& \neg \{ \neg [P = C] \& [P \mapsto (y \& z)] \} \qquad \qquad \qquad 4,Th.F1,lt,5,D.0.4
\end{aligned}$$


L. 4 derives from L. 3 by the single change of uncoupling the two lines of 'n' of which 'C' is the end point. L. 6: All the lines of 'j' are "conjoined" to form a single fundamental line by Th. F1 iterated, and similarly with 'k'. If after any addition the lines of 'j' or 'k' a line from one had an end-point in common with a line from the other, which point was not 'C' then, applying Th. F1, D. 0.4 would be controverted, hence the separation of the two sets.




The structure of a seriate set of points




Theorem 1.1




Theorem 1.2




Theorem 1.6




Theorem 1.9




Theorem 1.8




Theorem 2.3




Theorem 2.4




Theorem 2.5






Theorem 2.8



Theorem 2.11



Theorem 2.9



Theorem 2.14







Illustration of $\underline{p^M p^N} \mapsto t$



Line 9



Line 12



Line 10

A five country map